

When should hands be washed?

- Whenever the skin comes into contact with other people or with shared surfaces or equipment;
- Before and after athletic practice, games, or working out;
- After sneezing, coughing, blowing or touching the nose;
- Before and after touching the eyes, nose, or mouth, or skin that has open sores, boils and skin rashes;
- After using the toilet or urinal;
- After touching any item soiled with wound fluids, such as bandages, dressings or bedding;
- After cleaning the bathroom, changing your bedding, or doing laundry; and
- Before preparing food, eating, or drinking.



What else can be done to help prevent MRSA and staph infections?

- Bathe regularly,
- Take good care of the skin; avoid skin damage, clean damaged skin promptly with soap and water,
- Keep cuts and scrapes clean and covered with a bandage until healed.

Pus from infected wounds can contain staph and MRSA.

- Don't share personal items such as towels, clothing, or other items that touch skin, and use a barrier (e.g., clothing or a towel) between your skin and shared equipment such as weight-training benches.
- Wash soiled clothes, sheets, and towels with water and laundry detergent and dry completely in a hot dryer.



For more information about MRSA and staph infections, contact your doctor, local health department or school nurse.

References:

California Department of Public Health, MRSA for Athletes: What You Need to Know, <http://www.cdph.ca.gov/healthinfo/discond/Documents/CAMRSAForAthletes.pdf>

Centers for Disease Control and Prevention, Questions and Answers about Methicillin-Resistant *Staphylococcus aureus* (MRSA) in Schools,

<http://www.cdc.gov/Features/MRSAinSchools/>



A Guide to MRSA in California

What You Need to Know

Adapted from
"A Parent's Guide to MRSA in California" a
publication of the
California Department of Public Health
Division of Communicable Disease Control
Infectious Diseases Branch
www.cdph.ca.gov

What is Staphylococcus aureus?

Staphylococcus aureus (staph) are bacteria that many healthy people carry on their skin or in their nose. About 25% to 30% of



people in the United States carry staph in their nose, but it does not make them sick. Staph can also be carried in the armpit, groin, rectum or genital area. Most staph infections are minor and can be treated

without antibiotics. However, staph can sometimes cause serious infections like pneumonia, blood or joint infections, and deep skin infections.

What is MRSA?

Methicillin-resistant Staphylococcus aureus (MRSA) is a type of staph that is not killed by penicillin and similar antibiotics. About 1 out of every 100 people carries MRSA without making them sick. In California over half of all the staph infections are caused by MRSA. MRSA infections do not look any different than those caused by ordinary staph.

What do staph and MRSA infections look like?

Common skin conditions caused by staph and MRSA may look like any of the following:

Sores that look and feel like spider bites

(but are not spider bites);

Red painful bumps

under the skin, called

boils or abscesses;

A cut that is swollen,

hot, and filled with

pus;

Blisters filled with fluid or red skin with a

honey-colored crust (usually on the face); or

Red, warm, firm skin area that is painful and

getting larger (usually on the legs).



How does a person get staph or MRSA?

Staph and MRSA can be spread when a

person:

has direct skin-

to-skin contact

with an infected

person,

touch surfaces

that have staph

on them from

someone else's infection (e.g., towels,

athletic equipment, used bandages).

Staph is not usually passed through the air.



How are staph and MRSA infections treated?

Some staph skin infections are treated by

draining the sores and may not require

antibiotics. Draining these sores should be

done by a doctor.

If the doctor gives you antibiotics, make

sure you take all of the medicine, even if the

infection is getting better. Do not share your

medicine with anyone else or save it to use

at another time.

Call your doctor back if the infection does

not get better after a few days. If other

people you know or live with get the same

infection tell them to go to their doctor.

Is it possible to have another staph or

MRSA skin infection after it is cured?

Yes. It is possible to have another staph or

MRSA skin infection after it is treated. To

keep this from happening, follow the

doctor's orders while you have the infection,

and follow the prevention steps in this

pamphlet.

What should I do if I think my child has

MRSA?

If you think your child has MRSA, call a

doctor. Do not ignore the sore and hope it

will go away.

What is the easiest way to prevent MRSA

and staph infections?

Hand washing is the easiest way to prevent

MRSA and staph infections.

If soap and running water is not available

and hands do not have visible dirt on them,

staph germs can be killed by using an

alcohol-based hand rub.