

STANDARD PRECAUTIONS+

CONTACT ISOLATION

Staff must:



CLEAN HANDS*

*USE SOAP + WATER ON EXIT, IF APPROPRIATE (SEE REVERSE FOR DETAILS)



GOWN



GLOVES



USE
DISPOSABLE
EQUIPMENT
WHEN POSSIBLE

After discharge:



DISINFECT USED SURFACES

Ambulatory – CONTACT ISOLATION

Use for patients exhibiting diarrhea with incontinence, wound with uncontrolled drainage, lice, scabies, or other conditions per ISOLATION TABLE: https://infectioncontrol.ucsfmedicalcenter.org/isolation-table

D			CE		П		
Р	ĸ	U	CE	v	u	ĸ	E

SUPPLIES NEEDED

- THIS SIGN- hang at entryway to patient care area
- GOWN
- GLOVES

PATIENT

- No personal protective equipment (PPE) required
- Keep cover over wounds unless instructed to remove by provider
- Family members/caregivers accompanying patients do not need to wear PPE
- Clean hands prior to donning personal protective equipment (PPE)
- Wear gown and gloves upon entry to patient care area
- PPE Removal- Remove PPE before exiting patient care area
 - Grasp PPE in a manner that avoids contamination (Outside of PPE is contaminated)

STAFF

- Remove and discard disposable PPE
- Clean hands after removal of PPE (use soap and water for patients with active stooling, known or suspected C. difficile or Norovirus)
- Use dedicated disposable equipment (e.g. stethoscope, blood pressure cuff, thermometer), when possible
- Clean non-disposable equipment with facility-approved disinfecting wipe after each use

ROOM CLEANING

- Don and doff PPE as above
- Clean used surfaces and equipment with facility-approved disinfectant per standard cleaning procedures
- Change curtains when visibly soiled
- "High Clean" not required
- Remove isolation sign when cleaning complete, after discharge

For additional information, indications for, and discontinuation of isolation, refer to "Standard and Transmission-based Isolation Policy" and "Isolation Table" at http://infectioncontrol.ucsfmedicalcenter.org or contact Infection Prevention at 415-353-4343.

