

Measles Pediatric Quick Tips

There is a measles outbreak in Texas and New Mexico. In the last 2 months, there have been 5 measles cases in California.

3/13/25

Identify Consider measles	 In patients with fever, cough, coryza (runny nose), or conjunctivitis and a rash Maculopapular rash usually appears 3-5 days after symptoms begin, starts at hairline and descends. who, in the last 21 days, have traveled internationally or domestically to a region with a known measles outbreak including Texas and New Mexico. who are unvaccinated or have only received 1 MMR vaccine. who have been exposed to measles.
-------------------------------------	--

Refer to this clinical algorithm to help evaluate patients with fever/rash



Isolate Patients with suspected/ confirmed measles immediately	 -Mask patients immediately. For pediatric patients that cannot tolerate being masked, place a blanket loosely over the face if clinically safe. -Room the patient immediately -Prioritize placement of the patient into an Airborne Infection Isolation Room (AIIR) with the door closed. -In Ambulatory settings, place patient in a private exam room and close the door. Keep the door closed for the entire visit. -Staff entering the room should wear a fit-tested N95 or PAPR.

	BCH-San Francisco Business Hours, Mon-Fri 8am-4pm (628) 248-8503(Pediatric Voalte) (415)353-4343 (Main Line) After Hours, Hospital Supervisor (415)502-0728		MMR vaccine
Inform Call Infection Prevention right away	BCH-Oakland Call the pediatric ID physician on call Contact the Department of Public Health to	J. C. C. M.	remains the most important tool for preventing
	approve testing San Francisco County: (628)217-6100 Alameda County: Business Hours (510)268-3250; After Hours (925)422-7595		measles