

Clean your hands before and after touching PPE

Version 3.24.20

Extended use = wearing PPE continuously without doffing between multiple patient interactions with respiratory/droplet isolation (doff for interactions with patients not on isolation) Extended use is for specialty care respiratory units (15L, ED, RSCs, ACUs)

Reuse = doffing PPE and storing in a clean dry place after being cleaned per the guidance below Reuse is for ALL Clinical areas

For both Extended Use and Reuse, refer to Additional Nursing Practice Update Guidance for PPE use with Hazardous Medications

Regular Surgical Masks & Disposable N95 Respirators

Extended Use Guidance (15L, ED, RSCs, ACUs)

- Clean your hands each time prior to and after touching the mask/respirator
- **Wear a face shield with mask/respirator** for extended use (face shield will partially cover the mask)
- When exiting patient room, doff appropriate PPE (gown and gloves) and then clean hands
- Do not doff the mask/respirator when exiting a patient room
- Perform hand hygiene and don appropriate PPE as required (gloves/gown) before entering new patient room for a separate patient interaction

Extended Use Surgical Masks and/or Respirators can be reused after being doffed per the reuse guidance

Extended Use Guidance (15L, ED, RSCs, ACUs)

- Use one shield or goggles per health care worker
- Label with name and date
- When exiting patient room doff gown and gloves and then clean hands
- Do not doff the face shield, goggles or PAPR when exiting a patient room
- Perform hand hygiene and don appropriate PPE as required (gloves/gown) before entering new patient room for a separate patient interaction

Extended Use Face Shields, goggles, or PAPR shields can be reused after being doffed per the reuse guidance

Reuse Guidance

- Clean your hands each time prior to and after touching the mask/respirator
- **Wear a face shield with mask/respirator** (see pictures)
- Use your mask/respirator for a full shift, or longer if it does not meet discard guidance
- Do not share your mask/respirator
- When doffing, avoid touching the front of the mask/respirator
 - remove using elastic bands, loops or ties
- **Surgical Mask:** When donning for reuse avoid touching the front of the mask; don using elastic bands
- **Respirator:** When donning for reuse, clean hands immediately after touching the front of the respirator for fit check
- Avoid touching the inside of the mask/respirator unless hands are clean
- Between use: store in a clean, dry location that is labeled

Face Shields, PAPR Shields & Goggles

Reuse Guidance

- Clean your hands each time prior to putting on and taking off
- Use one shield or one pair of goggles, per health care worker
- Label with name and date; maintain control of your shield/goggles
- Clean external surfaces of the shield/goggles with hospital approved disinfectant wipe, followed by alcohol wipe to remove any residue if needed
 - PAPR Shield: Clean the external surface of the protective film that covers health care worker's chin
- Store shield/goggle in a clean, dry location when not being used.
 - PAPR shields can be left on PAPR and placed on a clean surface if the PAPR will be used by the same health care worker
- Relabel as needed if label was compromised during cleaning

Discard Guidance

Discard the surgical mask or respirator if:

- Contaminated hands have touched inside
- Wet, soiled, or damaged
- Appropriate fit is not maintained

Discard Guidance

Discard the face shield, goggles, or PAPR shield if:

- Inside cannot be cleaned
- Damaged in any manner
 - foam batting damaged/soiled
 - face shield cracked/ripped
 - Chin cuff of PAPR shield not snug to skin



Surgical mask with face shield



N-95 respirator with face shield



PAPR (CAPR) with PAPR shield