

GENERAL FAQs: COVID-19 Testing for Adult and Pediatric Patients

Is routine COVID-19 testing for all patients without COVID-19 symptoms still required on hospital admission and on hospital day 3?

- As of May 6, routine testing of all asymptomatic patients on hospital admission and on hospital day 3 is no longer required.
- See below for specific situations where testing of asymptomatic patients is still needed.
- In addition, providers can still order COVID-19 tests for asymptomatic patients based on clinical discretion.

Which asymptomatic patients still require testing?

- Patients with known, recent (i.e., within the previous 10 days) exposure to another person with confirmed COVID-19.
 - Note: Asymptomatic COVID-19 exposed patients should be tested for COVID-19 on the schedules recommended in the UCSF Health COVID-19 Quarantine Guidance.
 - Even if a patient has had COVID-19 in the last 90 days, they should still be tested for COVID-19 if they have had an exposure or develop symptoms.
- Specific patients may require COVID-19 testing for discontinuation of isolation precautions, including:
 - Select severely immunocompromised patients)
 - To enable earlier discontinuation of isolation for immunocompromised patients and patients who have had severe COVID-19 infection.
 - See UCSF Guidance for Discontinuation of Isolation for COVID-19 for details.
- Some inpatients may also require testing before discharge to congregate living situations

Testing indications	Recommendation
Asymptomatic	No longer required on admission or hospital day #3
Recent exposure to COVID-19	Refer to <u>quarantine guidance</u>
Symptoms consistent with COVID-19	Yes
Repeat testing for discontinuation of Novel	See UCSF guidance for discontinuation of isolation
Respiratory Isolation for high risk patients	See dest gardance for discontinuation of isolation

When should patients be tested for COVID-19?

 Any inpatient who develops signs or symptoms consistent with COVID-19 should be placed into Novel Respiratory Isolation and tested as a symptomatic patient. Patients with COVID-19 exposures should also be tested even if they are asymptomatic as per the <u>quarantine guidance</u>.

What personal protective equipment (PPE) is required when obtaining a COVID-19 specimen for testing?

• N95 respirator plus eye protection (or PAPR), gown, and gloves

What type of test is acceptable?

 Polymerase chain reaction (PCR) or other nucleic acid amplification test (NAAT), including the Abbott ID Now NAAT test.