

UCSF Health Interim COVID-19 Guidance for Symptomatic Adult ED, Inpatient OB Triage, Birth Center, and Hospitalized Patients

Guidelines are intended to assist with clinical decision-making for common situations but cannot replace personalized evaluation and management decisions based on individual patient factors


Isolation types:
-Droplet isolation: Surgical mask plus eye protection. "Droplet Isolation" sign placed on door
-Novel Respiratory Isolation: Contact + either N95 with eye protection or PAPR. "Novel Respiratory Isolation" sign placed on door

Signs available at: <https://infectioncontrol.ucsfmedicalcenter.org/ucsf>

UCSF Health V.8
6.21.2021
UCSF COVID-19 Care Delivery Committee Owners: Sarah Doernberg and Matt Aldrich

Adult ID services
COVID ID/MB/MtZ Attending
 Via Careweb or 415-443-0190
Transplant ID
 Via Careweb or 415-443-2552
Jacobs ID Attending
 Via Careweb or 415-443-8996

Hospital Epidemiology and Infection Prevention (HEIP)
Moffitt-Long and Mission Bay:
Business Hours
 Adult IC Practitioner on-call VOALTE: (628)248-9059
After Hours
Moffitt-Long:
 ML Hospital Supervisor Spectralink 415-353-8036 or 415-353-1964
Mission Bay Adults:
 MB Adult Hospital Clinical Resource Nurse 415-502-0562



Recommended testing:
 Single swab for OP plus either NP or Mid-turbinate
Acceptable testing materials:
 1. Flocked swab Nasopharyngeal (NP) plus oropharyngeal (OP)
 2. Flocked swab Mid-Turbinate plus OP
 3. Synthetic Swab Mid-Turbinate plus OP
-All tests can be requested through the same Apex order for COVID-19 RNA
-RVP and COVID-19 RNA testing should be done on the same swab
-Order either RVP or rapid flu/RSV but not both

Additional testing may be indicated based on clinical presentation

Transfer to a negative-pressure room and order "Airborne Isolation" at any time a patient requires intubation, high-flow nasal cannula, non-invasive ventilation, or tracheostomy

Refer to "Discontinuing Isolation for Patients with COVID-19" for criteria and process for isolation discontinuation

*Do not re-test asymptomatic patients who have "COVID recovered" flag in place

***Aerosol-generating Procedure Guidance**
https://infectioncontrol.ucsfmedicalcenter.org/sites/g/files/tksra4681/f/Guidance_for_PPE_use_for_High_Risk_Aerosol_Generating_Procedures.pdf
Surgical Procedure Guidance
https://infectioncontrol.ucsfmedicalcenter.org/sites/g/files/tksra4681/f/Asymptomatic_COVID_PPE.pdf

Isolation table
<https://infectioncontrol.ucsfmedicalcenter.org/isolation-table>